

A NOTE ON THE POINTS WITH DENSE ORBIT UNDER $\times 2$ AND $\times 3$ MAPS

AUTHOR ONE, AUTHOR TWO, AND AUTHOR THREE

ABSTRACT. It was conjectured by Furstenberg that for any $x \in [0, 1] \setminus \mathbb{Q}$,

$$\dim_H \overline{\{2^n x \pmod{1} : n \geq 1\}} + \dim_H \overline{\{3^n x \pmod{1} : n \geq 1\}} \geq 1.$$

When x is a normal number, the above result holds trivially. In this note, we give explicit non-normal numbers for which the above dimensional formula holds.

1. Introduction

Let $b \geq 2$ be an integer and T_b be the b -ary expansion given by

$$Tx = bx \pmod{1}.$$

x is a real number given by its b -ary expansion, that is,

$$x = [x] + \sum_{k \geq 1} \frac{a_k}{b^k} = [x] + 0.a_1 a_2 \cdots$$

where the digit $a_1, a_2, \dots$ are integers from $\{0, 1, \dots, b-1\}$ and an infinity of the a_k are not equal to $b-1$. Set

$$A_b(d, N, x) := \text{Card}\{j : 1 \leq j \leq N, a_j = d\}.$$

More generally, for a block $D_k = d_1 \cdots d_k$ of k digits from $\{0, 1, \dots, b-1\}$, set

$$A_b(D_k, N, x) := \text{Card}\{j : 0 \leq j \leq N-k, a_{j+1} = d_1, \dots, a_{j+k} = d_k\}$$

A real number x is normal to base b , if and only if, for every $k \geq 1$, every block of k digits from $\{0, 1, \dots, b-1\}$ occurs in the b -ary expansion of x with the same frequency $\frac{1}{b^k}$, that is, if and only if,

$$\lim_{n \rightarrow \infty} \frac{A_b(D_k, N, x)}{N} = \frac{1}{b^k}$$

for every $k \geq 1$ and every block D_k of k digits from $\{0, 1, \dots, b-1\}$.

Given one integer $b \geq 2$, the dynamical properties of the orbit $T_b^n(x)$ were well studied in the literature and are clearly known at least for almost all x . However, as far as two integers b_1 and b_2 are concerned (assume that they are multiplicatively independent), the properties of the orbits $T_{b_1}^x(x)$ and $T_{b_1}^n(x)$ with a common point x is far from being well understood. Very little is known on this topic. Intuitively, if

Date: December 26, 1997.

2000 Mathematics Subject Classification. Primary 05C38, 15A15; Secondary 05A15, 15A18.

Key words and phrases. Hausdorff dimension, dense of the orbit.

Thanks for Author One.

Thanks for Author Two.

This paper is in final form and no version of it will be submitted for publication elsewhere.

the expansion of x under T_{b_1} is simple, it cannot be too simple under T_{b_2} (for some evidence see [3]). And the normality of a number is closely related to his expansion to different bases([2],[?]), nowadays, the investigate of normal number has been used to resolve the problems in dynamical system and ergodic.

Furstenberg conjectured that [1]: for any $x \in [0, 1] \setminus \mathbb{Q}$,

$$\dim_H \overline{\{2^n x \pmod{1} : n \geq 1\}} + \dim_H \overline{\{3^n x \pmod{1} : n \geq 1\}} \geq 1.$$

It is clear that this is true for almost all x , since almost all x are normal numbers. So, a natural question is to ask, besides normal numbers, can one give explicit examples fulfilling the above conjecture. In this short note, we proof there exist a set E of full dimension which make Furstenberg conjectured hold.

2. Some notation

We use C_1 and C_2 to denote the collection of finite 2-adic and 3-adic words respectively, i.e.

$$C_1 = (\bigcup_{n \geq 1} \{0, 1\}^n), \quad C_2 = (\bigcup_{n \geq 1} \{0, 1, 2\}^n).$$

Let $\mathcal{A} = (w_1, v_1, w_2, v_2, \dots)$, where $w_i \in C_1$ and $v_i \in C_2$, be an enumeration of the words in C_1 and C_2 such that each word in C_1 and C_2 appears infinitely many times in $\mathcal{A}$.

Let

$$T_2 x = 2x \pmod{1} \text{ and } T_3 x = 3x \pmod{1}, x \in [0, 1]$$

be the $\times 2$ and $\times 3$ maps.

- $|w|$: the length of the word w ;
- $\lfloor \xi \rfloor$: the integer part of a real number ξ ;
- 0^m : a word of length m composed by 0.
- r_i : the integer $\lfloor |w_i| + (2 + |v_i|) \log_2 3 \rfloor + 2$;

For $u \in C_1$ and $v \in C_2$, the 2-adic cylinder and 3-adic cylinder are defined as

$$\begin{aligned} [u]_2 &:= \{x \in [0, 1] : \text{the 2-adic expansion of } x \text{ starts with } u\}, \\ [v]_3 &:= \{x \in [0, 1] : \text{the 3-adic expansion of } x \text{ starts with } v\}. \end{aligned}$$

3. Main results

Theorem 1. *There exist a Cantor set E composed of nonnormal numbers such that for all $x \in E$,*

- 1: $\overline{\{2^n x \pmod{1} : n \geq 1\}} = [0, 1]$ and $\overline{\{3^n x \pmod{1} : n \geq 1\}} = [0, 1]$.
- 2: $\dim_H E = 1$.

The construction of the Cantor set E is divided into three steps. Recall that $\mathcal{A} = \{w_1, v_1, w_2, v_2, \dots\}$ be an enumeration of the elements in $w \in C_1$ and $v \in C_2$ such that each element in $C_1 \cup C_2$ appears infinitely many times.

Fix an integer $m \geq 1$. Choose a sequence of integers $\{\ell_k\}_{k \geq 1}, \{n_k\}_{k \geq 1}$ such that

$$(3.1) \quad n_k = \ell_k \cdot m, \quad \lim_{k \rightarrow \infty} \frac{r_1 + \dots + r_k}{n_k} = 0.$$

For each $k \geq 1$, write

$$(3.2) \quad \widetilde{C}_1^{n_k} := \{(i_1, \dots, i_{n_k}) \in C_1 : i_{\ell_{m+1}} = i_{\ell_{m+m}} = 1, 0 \leq \ell < \ell_k\}.$$

Step 1: the first level of the Cantor set.

For each $(i_1^{(1)} \dots i_{n_1}^{(1)}) \in \widetilde{C}_1^{n_1}$, consider the 2-adic cylinder

$$[i_1^{(1)} \dots i_{n_1}^{(1)}, w_1]_2.$$

Let r_1 be the integer such that

$$3 \left(\frac{1}{3} \right)^{r_1} \leq \left(\frac{1}{2} \right)^{n_1 + |w_1|} < 3 \left(\frac{1}{3} \right)^{r_1 - 1}.$$

Then the 2-adic cylinder $[i_1^{(1)} \dots i_{n_1}^{(1)}, w_1]_2$ will contain at least one 3-adic cylinder of order r_1 . Denote by $[\epsilon_1]_3$ such a 3-adic cylinder (if there are many, just choose one).

It should be mentioned that r_1 does not depend on $(i_1^{(1)} \dots i_{n_1}^{(1)})$ but ϵ_1 does. This dependence will not play a role in the following argument, thus will not be explicitly addressed.

Now consider the 3-adic subcylinder $[\epsilon_1, v_1]_3$ contained in $[\epsilon_1]_3$. Similarly, let t_1 be the integer such that

$$2 \left(\frac{1}{2} \right)^{t_1} \leq \left(\frac{1}{3} \right)^{r_1 + |v_1|} < 2 \left(\frac{1}{2} \right)^{t_1 - 1}.$$

Then choose a 2-adic cylinder $[\eta_1]_2$ of order t_1 contained in $[\epsilon_1, v_1]_3$, so contained in the 2-adic cylinder $[i_1^{(1)} \dots i_{n_1}^{(1)}, w_1]_2$. Thus we can write this 2-adic cylinder $[\eta_1]_2$ as

$$[\eta_1]_2 = [i_1^{(1)} \dots i_{n_1}^{(1)}, w_1, \tilde{v}_1]_2,$$

to emphasize its dependence on v_1 . Now we have the following inclusions:

$$(3.3) \quad [i_1^{(1)} \dots i_{n_1}^{(1)}, w_1, \tilde{v}_1]_2 \subset [\epsilon_1, v_1]_3 \subset [i_1^{(1)} \dots i_{n_1}^{(1)}, w_1]_2.$$

A simple calculation can give us an estimation on the integer t_1 :

$$n_1 + |w_1| + (|v_1| + 1) \frac{\log 3}{\log 2} + 1 \leq t_1 < n_1 + |w_1| + (|v_1| + 2) \frac{\log 3}{\log 2} + 2.$$

The first level of E is defined as

$$F_1 = \bigcup_{(i_1^{(1)} \dots i_{n_1}^{(1)}) \in \widetilde{C}_1^{n_1}} [i_1^{(1)} \dots i_{n_1}^{(1)}, w_1, \tilde{v}_1]_2,$$

which is a collection of 2-adic cylinders of order t_1 .

Step k: the k -level of the Cantor set.

This is an inductive step. Assume that the $(k-1)$ th level F_{k-1} has been constructed, which is a collection of 2-adic cylinders $[\eta_{k-1}]$ of order t_{k-1} . Now we construct the k th level F_k .

Fix an element $[\eta_{k-1}]_2 \in F_{k-1}$. For each $(i_1^{(k)} \dots i_{n_k}^{(k)}) \in \widetilde{C}_1^{n_k}$, consider the 2-adic cylinder

$$[\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k]_2.$$

Similar as in Step 1, let r_k be the integer such that

$$3 \left(\frac{1}{3} \right)^{r_k} \leq \left(\frac{1}{2} \right)^{t_{k-1} + |w_k|} < 3 \left(\frac{1}{3} \right)^{r_k - 1}.$$

Then there is a 3-adic cylinder, denoted by $[\epsilon_k]_3$, of order r_k contained in the 2-adic cylinder $[\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k]_2$ (if there are many, just choose one).

It should also be mentioned that r_k does not depend on $(i_1^{(j)} \dots i_{n_j}^{(j)})$ for $1 \leq j \leq k$ but ϵ_k does. This dependence will not play a role in the following argument, thus will not be explicitly addressed.

Now consider the 3-adic subcylinder $[\epsilon_k, v_k]_3$ contained in $[\epsilon_k]_3$. Let t_k be the integer such that

$$2 \left(\frac{1}{2} \right)^{t_k} \leq \left(\frac{1}{3} \right)^{r_k + |v_k|} < 2 \left(\frac{1}{2} \right)^{t_k - 1}.$$

Then choose a 2-adic cylinder $[\eta_2]_2$ of order t_k contained in $[\epsilon_k, v_k]_3$, so contained in the 2-adic cylinder $[\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k]_2$. Thus we can write this 2-adic cylinder $[\eta_k]_2$ as

$$[\eta_k]_2 = [\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k, \tilde{v}_k]_2,$$

to emphasize its dependence on v_k . Now we have the following inclusions:

$$(3.4) \quad [\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k, \tilde{v}_k]_2 \subset [\epsilon_k, v_k]_3 \subset [\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k]_2.$$

A simple calculation can give us an estimation on the integer t_k (replace the role of n_1 by t_{k-1} , $|w_1|$ by $|w_k|$, $|v_1|$ by $|v_k|$):

$$(3.5) \quad t_{k-1} + |w_k| + n_k + (|v_k| + 1) \frac{\log 3}{\log 2} + 1 \leq t_k < t_{k-1} + |w_k| + n_k + (|v_k| + 2) \frac{\log 3}{\log 2} + 2.$$

Let $p_i = |w_k, \tilde{v}_k| = t_k - t_{k-1} - n_k \leq r_i$. Thus it follows from (3.5) and (3.1) that

$$(3.6) \quad \lim_{k \rightarrow \infty} \frac{|w_1, \tilde{v}_1| + \dots + |w_k, \tilde{v}_k|}{n_1 + \dots + n_k} \leq \lim_{k \rightarrow \infty} \frac{r_1 + \dots + r_k}{n_1 + \dots + n_k} = 0.$$

The k th level of E is defined as

$$F_k = \bigcup_{[\eta_{k-1}]_2 \in F_{k-1}} \bigcup_{(i_1^{(k)} \dots i_{n_k}^{(k)}) \in \widetilde{C}_1^{n_k}} [\eta_{k-1}, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k, \tilde{v}_k]_2,$$

which is a collection of 2-adic cylinders of order t_k .

Finally, the desired Cantor set E is defined as

$$E = \bigcup_{m \geq 1} E_m, \quad E_m = \bigcap_{k \geq 1} \bigcup_{[\eta_k]_2 \in F_k} [\eta_k]_2.$$

The following three propositions correspond to the three items in Theorem 1.

Proposition 1. *Every point in E_m is nonnormal.*

Proof. By the construction of E_m , for each $x \in E_m$, we express x by its 2-adic expansion as follows:

$$x = [i_1^{(1)} \dots i_{n_1}^{(1)}, w_1, \tilde{v}_1, \dots, i_1^{(k)} \dots i_{n_k}^{(k)}, w_k, \tilde{v}_k, \dots]_2,$$

where $(i_1^{(k)} \dots i_{n_k}^{(k)}) \in \widetilde{C}_1^{n_k}$.

We count the frequency of the word 0^{2^m} occurring in the 2-adic expansion of x . Recall the definition of $\widetilde{C}_1^{n_k}$. Among the first $t_{k-1} + n_k$ terms in the 2-adic expansion of x , the number of the occurrence of 0^m will be less than $p_1 + \dots + p_{k-1}$. Thus the frequency of the word 0^m occurring in the 2-adic expansion of x is less than

$$\liminf_{k \rightarrow \infty} \frac{p_1 + \dots + p_{k-1}}{n_k} = 0.$$

This shows that x is nonnormal.

This shows that x is nonnormal. $\square$

Proposition 2. *For any $x \in E$,*

$$\overline{\{2^n x \pmod{1} : n \geq 1\}} = [0, 1], \quad \overline{\{3^n x \pmod{1} : n \geq 1\}} = [0, 1].$$

Proof. Fix an $x \in E$. It is sufficient to show that for any 2-adic finite word $w \in C_1$ and $v \in C_2$, they appear infinitely often in the 2-adic expansion and the 3-adic expansion of x respectively.

Recall the definition of $\mathcal{A}$. It suffices to show that for each $k \geq 1$, w_k and v_k appear in the 2-adic expansion and the 3-adic expansion of x respectively. This follows from (3.4) by noticing that

$$T_3^{r_k} x \in [v_k]_3, \quad T_2^{t_{k-1} + n_k}(x) \in [w_k]_2.$$

□

Proposition 3. *For each $m \geq 2$, $\dim_H E_m \geq 1 - 2/m$.*

Proof. Recall the definition of $\widetilde{C}_1^{n_k}$ (3.2). Write the points in E_m by its 2-adic expansion:

$$\left\{ [i_1^{(1)} \cdots i_{n_1}^{(1)}, w_1, \tilde{v}_1, \dots, i_1^{(k)} \cdots i_{n_k}^{(k)}, w_k, \tilde{v}_k, \dots]_2, \quad (i_1^{(k)} \cdots i_{n_k}^{(k)}) \in \widetilde{C}_1^{n_k}, k \geq 1 \right\}.$$

Now we will distribute a measure uniformly distributed on the cylinders with nonempty intersection with E_m . Thus we count the number of t -th cylinders which have nonempty intersection with E_m for each $t \geq 1$. We denote this number by b_t .

Recall the construction of the k th level F_k of the Cantor set E_m . We know that among the digit sequence of a point in E_m , at the positions

$$\left\{ t_k + \ell m + 1, t_k + (\ell + 1)m, t_k + n_{k+1} + 1, \dots, t_k + n_{k+1} + p_{k+1}, 0 \leq \ell < \ell_{k+1}, k \geq 0 \right\},$$

the digits are fixed, while at other positions, the digits can be chosen arbitrarily in $\{0, 1\}$. Thus the numbers b_t can be given as follows.

(i). When $t \leq t_1 = n_1 + p_1 = \ell_1 m + p_1$,

$$b_t = \begin{cases} 2^{t-2\ell}, & \text{when } t = \ell m \text{ for } 0 \leq \ell < \ell_1; \\ 2^{t-2\ell-1}, & \text{when } \ell m < t < (\ell + 1)m \text{ for } 0 \leq \ell < \ell_1; \\ 2^{\ell_1 m - 2\ell_1}, & \text{when } \ell_1 m \leq t \leq n_1 + p_1. \end{cases}$$

(ii). When $t_k < t \leq t_{k+1} = t_k + n_{k+1} + p_{k+1} = t_k + \ell_{k+1} m + p_{k+1}$,

$$b_t = \begin{cases} \prod_{j=1}^k 2^{\ell_j m - 2\ell_j} \cdot 2^{t-t_k-2\ell}, & \text{when } t = t_k + \ell m \text{ for } 0 \leq \ell < \ell_{k+1}; \\ \prod_{j=1}^k 2^{\ell_j m - 2\ell_j} \cdot 2^{t-t_k-2\ell-1}, & \text{when } \ell m < t < (\ell + 1)m \text{ for } 0 \leq \ell < \ell_{k+1}; \\ \prod_{j=1}^{k+1} 2^{\ell_j m - 2\ell_j}, & \text{when } t_k + n_{k+1} \leq t \leq t_k + n_{k+1} + p_{k+1}. \end{cases}$$

Thus for each 2-adic cylinder I_t with nonempty intersection with E_m , its measure can be given explicitly as

$$\mu(I_t) = b_t^{-1}.$$

As a consequence, together with (3.6), we have

$$(3.7) \quad \liminf_{t \rightarrow \infty} \frac{\log \mu(I_t)}{\log |I_t|} = \liminf_{t \rightarrow \infty} \frac{\log b_t}{t \log 2} \geq 1 - \frac{2}{m}.$$

So, using Billingsley's lemma [?]: Let $A \subset [0, 1]$ be Borel and let μ be a finite Borel measure on $[0, 1]$. Suppose $\mu(A) \geq 0$. If

$$\alpha \leq \liminf_{n \rightarrow \infty} \frac{\log \mu(I_n(x))}{\log |I_n(x)|} \leq \beta$$

for all $x \in A$, then $\alpha \leq \dim(A) \leq \beta$. we conclude that

$$\dim_H E_m \geq 1 - \frac{2}{m}$$

and then $\dim_H E = 1$. □

REFERENCES

- [1] Harry Furstenberg. Intersections of cantor sets and transversality of semigroups *Sympos. Salmomon. Bochner, princeton univ, princeton, N. J.* , (1969), 41–59.
- [2] Peter Hertling, Simply normal numbers to different bases., *Journal of Univesal Computer Science*, **8(2)** (2002), 235-242(electronic).
- [3] Bugeaud, Yann, Distribution modulo one and diophantin approximation *Cambridge tracts in mathematics* , 2012

(A. One and A. Two) AUTHOR ONE TWO ADDRESS LINE 1, AUTHOR ONE TWO ADDRESS LINE 2
E-mail address, A. One: `aone@aoneinst.edu`
URL: `http://www.authorone.oneuniv.edu`

Current address, A. Two: Author Two current address line 1, Author Two current address line 2

E-mail address, A. Two: `atwo@atwoinst.edu`
URL: `http://www.authortwo.twouniv.edu`

(A. Three) AUTHOR THREE ADDRESS LINE 1, AUTHOR THREE ADDRESS LINE 2
URL: `http://www.authorthree.threeuniv.edu`